

approval. Thoracoscopic treatment of the LAA has been suggested as a sole therapy for patients with AF to minimize stroke risk.⁸ There is evidence to suggest that the AtriClip device results in electrical isolation, and the device has been used as an isolated therapy for focal atrial tachycardia.²¹

Although the device studied is third generation, the deployment tool was a first-generation instrument. This reusable large deployment tool was cumbersome and can be used only via a sternotomy. There are 2 currently approved deployment tools. One is designed for open sternotomy, and the other is used for thoracoscopic approach. Both are smaller and more flexible, and have a lower profile than the deployment tool used in this report.

Limitations

This study is limited in the short-term imaging follow-up of only 3 months, although clinical follow-up extends to 12 months. This is a relatively small cohort of patients. Longer follow-up is needed to evaluate for evidence of device migration. This study was not designed to assess reduction in stroke risk. Late neurologic events developed in 2 patients, which did not appear to be related to the LAA to the best of our knowledge. A significantly larger randomized study would be required with longer-term follow-up to document any efficacy in stroke prophylaxis.

CONCLUSIONS

This multicenter initial trial suggests that exclusion of the LAA can be performed safely and without injury to the heart or surrounding structures during open cardiac surgery with the AtriClip device. In short-term follow-up, there is evidence of persistent LAA closure with no communication to the appendage. Long-term studies should be performed to evaluate the efficacy in the prevention of stroke and rule out the potential for device migration.

References

1. Wolf PA, Kannel WB, McGee DL, Meeks SL, Bharucha NE, McNamara PM. Duration of atrial fibrillation and imminence of stroke: the Framingham study. *Stroke*. 1983;14:664-7.
2. Blackshear JL, Odell JA. Appendage obliteration to reduce stroke in cardiac surgical patients with atrial fibrillation. *Ann Thorac Surg*. 1996;61:755-9.
3. Arantes L, Matsuo S, Lim KT, Knecht S, Jais P, Haissaguerre MJ. Left atrial appendage activity mimicking an epsilon wave on the surface electrocardiogram. *Cardiovasc Electrophysiol*. 2007;18:1114-5.
4. Calkins H, Brugada J, Packer DL, Cappato R, Chen SA, Crijns HJ, et al. HRS/EHRA/ECAS expert consensus statement on catheter and surgical ablation of atrial fibrillation: recommendations for personnel, policy, procedures and follow-up. *Europace*. 2007;9:335-79.
5. Katz ES, Tsiamtsiouris T, Applebaum RM, Schwartzbard A, Tunick PA, Kronzon I. Surgical left atrial appendage ligation is frequently incomplete: a transesophageal echocardiographic study. *J Am Coll Cardiol*. 2000;36:468-71.
6. Kanderian AS, Gillinov AM, Pettersson GB, Blackstone E, Klein AL. Success of surgical left atrial appendage closure: assessment by transesophageal echocardiography. *J Am Coll Cardiol*. 2008;52:924-9.
7. Bruce CJ, Stanton CM, Asirvatham SJ, Danielsen AJ, Johnson SB, Packer DL, et al. Percutaneous epicardial left atrial appendage closure: intermediate-term results. *J Cardiovasc Electrophysiol*. 2011;22:64-70.
8. Blackshear JL, Johnson WD, Odell JA, Baker VS, Howard M, Pearce L, et al. Thoracoscopic extracardiac obliteration of the left atrial appendage for stroke risk reduction in atrial fibrillation. *J Am Coll Cardiol*. 2003;42:1249-52.
9. Fumoto H, Gillinov AM, Ootaki Y, Akiyama M, Saeed D, Horai T, et al. A novel device for left atrial appendage exclusion: the third-generation atrial exclusion device. *J Thorac Cardiovasc Surg*. 2008;136:1019-27.
10. Salzberg SP, Gillinov AM, Anyanwu A, Castillo J, Filsoufi F, Adams DH. Surgical left atrial appendage occlusion: evaluation of a novel device with magnetic resonance imaging. *Eur J Cardiothorac Surg*. 2008;34:766-70.
11. Salzberg SP, Plass A, Emmert MY, Desbiolles L, Alkadhi H, Grünenfelder J, et al. Left atrial appendage clip occlusion: early clinical results. *J Thorac Cardiovasc Surg*. 2010;139:1269-74.
12. Bleeding during antithrombotic therapy in patients with atrial fibrillation. The Stroke Prevention in Atrial Fibrillation Investigators. *Arch Intern Med*. 1996;156:409-16.
13. Healey JS, Crystal E, Lamy A, Teoh K, Semelhago L, Hohnloser SH, et al. Left Atrial Appendage Occlusion Study (LAAOS): results of a randomized controlled pilot study of left atrial appendage occlusion during coronary bypass surgery in patients at risk for stroke. *Am Heart J*. 2005;150:288-93.
14. García-Fernández MA, Pérez-David E, Quiles J, Peralta J, García-Rojas I, Bermejo J, et al. Role of left atrial appendage obliteration in stroke reduction in patients with mitral valve prosthesis: a transesophageal echocardiographic study. *J Am Coll Cardiol*. 2003;42:1253-8.
15. Rosenzweig BP, Katz E, Kort S, Schloss M, Kronzon I. Thromboembolism from a ligated left atrial appendage. *J Am Soc Echocardiogr*. 2001;14:396-8.
16. Gillinov AM, Pettersson G, Cosgrove DM. Stapled excision of the left atrial appendage. *J Thorac Cardiovasc Surg*. 2005;129:679-80.
17. Holmes DR, Reddy VY, Turi ZG, Doshi SK, Sievert H, Buchbinder M, et al. Percutaneous closure of the left atrial appendage versus warfarin therapy for prevention of stroke in patients with atrial fibrillation: a randomised non-inferiority trial. *Lancet*. 2009;374:534-42.
18. Lynch M, Shanewise JS, Chang GL, Martin RP, Clements SD. Recanalization of the left atrial appendage demonstrated by transesophageal echocardiography. *Ann Thorac Surg*. 1997;63:1774-5.
19. Singh SM, Dukkhipati SR, d'Avila A, Doshi SK, Reddy VY. Percutaneous left atrial appendage closure with an epicardial suture ligation approach: a prospective randomized pre-clinical feasibility study. *Heart Rhythm*. 2010;7:370-6.
20. Bartus K, Bednarek J, Myc J, Kapelak B, Sadowski J, Lelakowski J, et al. Feasibility of closed-chest ligation of the left atrial appendage in humans. *Heart Rhythm*. 2011;8:188-93.
21. Benussi S, Mazzone P, Maccabelli G, Vergara P, Grimaldi A, Pozzoli A, et al. Thoracoscopic appendage exclusion with an AtriClip device as a solo treatment for focal atrial tachycardia. *Circulation*. 2011;123:1575-8.
22. Gage BF, Waterman AD, Shannon W, Boechler M, Rich MW, Radford MJ. Validation of clinical classification schemes for predicting stroke: results from the National Registry of Atrial Fibrillation. *JAMA*. 2001;285:2864-70.
23. Jaber WA, White RD, Kuzmiak SA, Boyle JM, Natale A, Apperson-Hansen C, et al. Comparison of ability to identify left atrial thrombus by three dimensional tomography versus transesophageal echocardiography in patients with atrial fibrillation. *Am J Cardiol*. 2004;93:486-9.
24. Wazni OM, Tsao HM, Chen SA, Chuang HH, Saliba W, Natale A, et al. Cardiovascular imaging in the management of atrial fibrillation. *J Am Coll Cardiol*. 2006;48:2077-84.

Discussion

Dr Bryan Meyers (St Louis, Mo). There was a long list of inclusion and exclusion criteria. How restrictive was this trial and how representative is this patient population compared with the group of patients you encounter every day in your cardiac surgery practice?

Dr Ailawadi. On the basis of the exclusion criteria from this study, I would say approximately half of the patients I see in daily practice could have qualified for this. The most liberal of the inclusion criteria was age greater than 65 years and hypertension. So many of the patients we treat meet that criteria in this trial.

Dr Wiley Nifong (Greenville, NC). Great study, great results. We have been very happy using the clip clinically as well. I did

notice early in the learning curve it looked like there were 3 misfires. Of course, while you are using it, you can get it down seated well. It appears, in the beating heart, to guide itself down. After it fires, I assume you could just use a wire cutter? You could remove it if you placed it and noticed it was not seated correctly, is that correct?

Dr Ailawadi. There are a couple of points to mention about deploying the device. The deployment device we used in this study was a large metal reusable tool, with a manual release. Once you release it, it is difficult, but it can be removed. You do not need a wire cutter, but you essentially take 2 pick-ups, separate the clip, and pull it off and then reattach it to the deployment tool and put it on again. The current FDA-approved device has a mechanism to open and close, and once you are happy with positioning, then you go ahead and cut the 4 sutures. So the current deployment device allows repositioning until it is in optimal position.

Dr Azhar Hossain (*Indianapolis, Ind*). How do you determine that you have been successful in excluding the entire appendage? As you know, there is residual appendage left, and the risk of thrombosis is still present. So are you using TEE assistance during the procedure?

Dr Ailawadi. It certainly can be done. I can tell you that during a sternotomy when you have the heart lifted up, it can be challenging to visualize by TEE. So in this situation, clip placement is by visual inspection. We do use TEE routinely in all of our cases. I have also performed this multiple times via a minimally invasive approach when performing a hybrid AF procedure or thoracoscopic AF procedure. In this situation, the heart is in its native position, and we can visualize by TEE before we deploy the clip.

Dr Bryan Meyers (*St Louis, Mo*). So you showed efficacy in the ability to exclude the appendage. The next step is going to be to show clinical effectiveness, and you are going to need a huge trial, aren't you, for the rare events you are trying to prevent?

Dr Ailawadi. Correct.

Dr Meyers. Any idea of where that is going next?

Dr Ailawadi. This is the million dollar question. I suspect it will require a large trial of at least 2000 patients. We will need to see what type of funding is available for this type of study. Ultimately, this will be required, not just with this particular device but with all the devices to see the clinical effectiveness of stroke reduction.